

- Swallow capsule whole. Do not chew, break, or crush.
- This medicine is given as a shot into a muscle.
- Follow diet plan and exercise program as recommended by healthcare provider.

What do I do if I miss a dose? (does not apply to patients in the hospital)

- Take a missed dose as soon as possible.
- If it is almost time for the next dose, skip the missed dose and return to your regular schedule.
- Do not take a double dose or extra doses.
- Do not change dose or stop medicine. Talk with healthcare provider.

What are the precautions when taking this medicine?

- If you are 65 or older, use this medicine with caution. You could have more side effects.
- If you have breast cancer, talk with healthcare provider.
- If you have diabetes, talk with healthcare provider.
- If you have a family history of diabetes, talk with healthcare provider.
- If you have Parkinson's disease, talk with healthcare provider.
- If you have seizures, talk with healthcare provider.
- If you have a slow heartbeat and do not have a pacemaker, talk with healthcare provider.
- If you have a weakened heart, talk with healthcare provider.
- If you are overweight, talk with healthcare provider.
- Check medicines with healthcare provider. This medicine may not mix well with other medicines.
- You may not be alert. Avoid driving, doing other tasks or activities until you see how this medicine affects you.
- Avoid alcohol (includes wine, beer, and liquor) or other medicines and natural products that slow your actions and reactions. These include sedatives, tranquilizers, mood stabilizers, antihistamines, and other pain medicine.
- Tell healthcare provider if you are pregnant or plan on getting pregnant.

What are some possible side effects of this medicine?

- Feeling lightheaded, sleepy, having blurred vision, or a change in thinking clearly. Avoid driving, doing other tasks or activities that require you to be alert or have clear vision until you see how this medicine affects you.
- Feeling dizzy. Rise slowly over several minutes from sitting or lying position. Be careful climbing.
- Nervous and excitable.
- Nausea or vomiting. Small frequent meals, frequent mouth care, sucking hard, sugar-free candy, or chewing sugar-free gum may help.
- Constipation. More liquids, regular exercise, or a fiber-containing diet may help. Talk with healthcare provider about a stool softener or laxative.