

# Older drugs - 50s first antipsychotic

## *Thorazine side effects may include:*

Abnormal secretion of milk, abnormalities in movement and posture, agitation, anemia, asthma, blood disorders, breast development in males, chewing movements, constipation, difficulty breathing, difficulty swallowing, dizziness, drooling, drowsiness, dry mouth, ejaculation problems, eye problems causing fixed gaze, fainting, fever, flu-like symptoms, fluid accumulation and swelling, headache, heart attack, high or low blood sugar, hives, impotence, inability to urinate, inability to move or talk, increase of appetite, infections, insomnia, intestinal blockage, involuntary movements of arms and legs, tongue, face, mouth, or jaw, irregular blood pressure, pulse, and heartbeat, irregular or no menstrual periods, jitteriness, light-headedness (on standing up), lockjaw, mask-like face, muscle stiffness and rigidity, narrow or dilated pupils, nasal congestion, nausea, pain and stiffness in the neck, persistent, painful erections, pill-rolling motion, protruding tongue, puckering of the mouth, puffing of the cheeks, rapid heartbeat, red or purple spots on the skin, rigid arms, feet, head, and muscles (including the back), seizures, sensitivity to light, severe allergic reactions, shuffling walk, skin inflammation and peeling, sore throat, spasms in jaw, face, tongue, neck, mouth, and feet, sweating, swelling of breasts in women, swelling of the throat, tremors, twitching in the body, neck, shoulders and face, twisted neck, visual problems, weight gain, yellowed skin and whites of eyes